

Annexe 1: Supporting References for Each Section – For Online Access

Assessment Processes for Frailty

- Morley JE, Adams EV. Rapid Geriatric Assessment. Journal of the American Medical Directors Association. 2015;16(10):808-12.
- Hubbard RE, Andrew MK, Fallah N, Rockwood K. Comparison of the prognostic importance of diagnosed diabetes, co-morbidity and frailty in older people. Diabetic medicine : a journal of the British Diabetic Association. 2010;27(5):603-6.
- Castro-Rodriguez M, Carnicero JA, Garcia-Garcia FJ, Walter S, Morley JE, Rodriguez-Artalejo F, et al. Frailty as a Major Factor in the Increased Risk of Death and Disability in Older People With Diabetes. Journal of the American Medical Directors Association. 2016;17(10):949-55.
- Morley JE, Malmstrom TK, Rodriguez-Manas L, Sinclair AJ. Frailty, sarcopenia and diabetes. Journal of the American Medical Directors Association. 2014;15(12):853-9.
- Wang Y, Xu XY, Feng CH, Li YL, Ge X, Zong GL, et al. Patients with type 2 diabetes exhibit cognitive impairment with changes of metabolite concentration in the left hippocampus. Metabolic brain disease. 2015;30(4):1027-34.
- Monette MC, Baird A, Jackson DL. A meta-analysis of cognitive functioning in nondemented adults with type 2 diabetes mellitus. Canadian journal of diabetes. 2014;38(6):401-8.
- Abner EL, Nelson PT, Kryscio RJ, Schmitt FA, Fardo DW, Woltjer RL, et al. Diabetes is associated with cerebrovascular but not Alzheimer's disease neuropathology. Alzheimer's & dementia : the journal of the Alzheimer's Association. 2016;12(8):882-9.
- Morley JE. Anorexia and weight loss in older persons. The journals of gerontology Series A, Biological sciences and medical sciences. 2003;58(2):131-7.
- Rubenstein LZ, Josephson KR, Wieland GD, English PA, Sayre JA, Kane RL. Effectiveness of a geriatric evaluation unit. A randomized clinical trial. The New England journal of medicine. 1984;311(26):1664-70.
- Stuck AE, Siu AL, Wieland GD, Adams J, Rubenstein LZ. Comprehensive geriatric assessment: a meta-analysis of controlled trials. Lancet. 1993;342(8878):1032-6.
- Van Craen K, Braes T, Wellens N, Denhaerynck K, Flamaing J, Moons P, et al. The effectiveness of inpatient geriatric evaluation and management units: a systematic review and meta-analysis. Journal of the American Geriatrics Society. 2010;58(1):83-92.
- Craig C, Chadborn N, Sands G, Tuomainen H, Gladman J. Systematic review of EASY-care needs assessment for community-dwelling older people. Age and aging. 2015;44(4):559-65.
- Vellas B, Balardy L, Gillette-Guyonnet S, Abellan Van Kan G, Ghisolfi-Marque A, Subra J, et al. Looking for frailty in community-dwelling older persons: the Gerontopole Frailty Screening Tool (GFST). The journal of nutrition, health & aging. 2013;17(7):629-31.
- Fukutomi E, Okumiya K, Wada T, Sakamoto R, Ishimoto Y, Kimura Y, et al. Importance of cognitive assessment as part of the "Kihon Checklist" developed by the Japanese Ministry of Health, Labor and Welfare for prediction of frailty at a 2-year follow up. Geriatrics & gerontology international. 2013;13(3):654-62.
- Li Y, Zou Y, Wang S, Li J, Jing X, Yang M, et al. A Pilot Study of the FRAIL Scale on Predicting Outcomes in Chinese Elderly People With Type 2 Diabetes. Journal of the American Medical Directors Association. 2015;16(8):714 e7- e12.
- Woo J, Leung J, Morley JE. Validating the SARC-F: a suitable community screening tool for sarcopenia? Journal of the American Medical Directors Association. 2014;15(9):630-4.
- Woo J, Yu R, Wong M, Yeung F, Wong M, Lum C. Frailty Screening in the Community Using the FRAIL Scale. Journal of the American Medical Directors Association. 2015;16(5):412-9.
- Rejeski WJ, Bray GA, Chen SH, Clark JM, Evans M, Hill JO, et al. Aging and physical function in type 2 diabetes: 8 years of an intensive lifestyle intervention. The journals of gerontology Series A, Biological sciences and medical sciences. 2015;70(3):345-53.
- Ngandu T, Lehtisalo J, Solomon A, Levalahti E, Ahtiluoto S, Antikainen R, et al. A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): a randomised controlled trial. Lancet. 2015;385(9984):2255-63.
- Ng TP, Feng L, Nyunt MS, Feng L, Niti M, Tan BY, et al. Nutritional, Physical, Cognitive, and Combination Interventions and Frailty Reversal Among Older Adults: A Randomized Controlled Trial. The American journal of medicine. 2015;128(11):1225-36 e1.

Glucose Regulation

- Global Guidelines for Managing Older People with Type 2 Diabetes. 2014: Available from: <http://www.idf.org/guidelines-older-people-type-2-diabetes>.
- Bonds DE, Miller ME, Bergenstal RM, Buse JB, Byington RP, Cutler JA, et al. The association between symptomatic, severe hypoglycemia and mortality in type 2 diabetes: retrospective epidemiological analysis of the ACCORD study. BMJ. 2010;340:b4909.
- Abbatecola AM, Bo M, Armellini F, D'Amico F, Desideri G, Falaschi P, et al. Tighter glycemic control is associated with ADL physical dependency losses in older patients using sulfonylureas or metformin: Results from the DIMORA study. Metabolism. 2015 Nov;64(11):1500-6.
- Monami M, Vitale V, Lamanna C, Bartoli N, Martelli D, Zannoni S, et al. HbA1c levels and all-cause mortality in type 2 diabetic patients: epidemiological evidence of the need for personalised therapeutic targets. Nutr Metab Cardiovasc Dis. 2013 Apr;23(4):300-6.
- Munshi MN, Florez H, Huang ES, Kalyani RR, Mupanomunda M, Pandya N, et al. Management of Diabetes in Long-term Care and Skilled Nursing Facilities: A Position Statement of the American Diabetes Association. Diabetes Care. 2016 Feb;39(2):308-18.
- Munshi MN, Segal AR, Slyne C, Samur AA, Brooks KM, Horton ES. Shortfalls of the use of HbA1C-derived eAG in older adults with diabetes. Diabetes Res Clin Pract. 2015 Oct;110(1):60-5.
- Munshi MN, Slyne C, Segal AR, Saul N, Lyons C, Weinger K. Liberating A1C goals in older adults may not protect against the risk of hypoglycemia. J Diabetes Complications. 2017 Mar 14.
- Karyekar CS, Ravichandran S, Allen E, Fleming D, Frederick R. Tolerability and efficacy of glycemic control with saxagliptin in older patients (aged \geq 65 years) with inadequately controlled type 2 diabetes mellitus. Clin Interv Aging. 2013;8:419-30.
- Munshi M, Slyne C, Segal AR, Saul N, Lyons C, Weinger K. Simplification of Insulin Regimen in Older Adults and Risk of Hypoglycemia. JAMA Internal Medicine. [Research Letter]. 2016 June 6, 2016;Online First.

Sinclair A, Morley JE, Rodriguez-Manas L, Paolisso G, Bayer T, Zeyfang A, et al. Diabetes mellitus in older people: position statement on behalf of the International Association of Gerontology and Geriatrics (IAGG), the European Diabetes Working Party for Older People (EDWPOP), and the International Task Force of Experts in Diabetes. *J Am Med Dir Assoc.* 2012 Jul;13(6):497-502.

Sinclair AJ. Good clinical practice guidelines for care home residents with diabetes: an executive summary. *Diabet Med.* 2011 Jul;28(7):772-7.

Kirkman MS, Briscoe VJ, Clark N, Florez H, Haas LB, Halter JB, et al. Diabetes in older adults. *Diabetes Care.* 2012 Dec;35(12):2650-64.

Claesens M, Gillard P, De Smet F, Callens M, De Moor B, Mathieu C. Mortality in Individuals Treated With Glucose-Lowering Agents: A Large, Controlled Cohort Study. *J Clin Endocrinol Metab.* 2016 Feb;101(2):461-9.

Du YF, Ou HY, Beverly EA, Chiu CJ. Achieving glycemic control in elderly patients with type 2 diabetes: a critical comparison of current options. *Clin Interv Aging.* 2014;9:1963-80.

Inzucchi SE, Bergenfelz RM, Buse JB, Diamant M, Ferrannini E, Nauck M, et al. Management of hyperglycemia in type 2 diabetes, 2015: a patient-centered approach: update to a position statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care.* 2015 Jan;38(1):140-9.

Mooradian AD. Special considerations with insulin therapy in older adults with diabetes mellitus. *Drugs Aging.* 2011 Jun 01;28(6):429-38.

Munshi MN SA, Ryan C, Slyne C, Saul N. Simplification of insulin regimens in older adults with diabetes decreases hypoglycemia, without worsening glycemic control. Scientific sessions of the American Diabetes Association; June 2013; Chicago, IL 2013. p. 390P.

Bonds DE, Miller ME, Dull J, Feinglos M, Ismail-Beigi F, Malozowski S, et al. Severe hypoglycemia symptoms, antecedent behaviors, immediate consequences and association with glycemia medication usage: Secondary analysis of the ACCORD clinical trial data. *BMC Endocr Disord.* 2012 May 30;12:5.

Exercise Intervention

Park SW, Goodpaster BH, Strotmeyer ES, Kuller LH, Broudeau R, Kammerer C, de Rekeneire N, Harris TB, Schwartz AV, Tylavsky FA, Cho YW, Newman AB; Health, Aging, and Body Composition Study. Accelerated loss of skeletal muscle strength in older adults with type 2 diabetes. *Diabetes Care.* 2007; 30(6):1507-1512

Volpatto S, Bianchi L, Lauretani F, Lauretani F, Bandinelli S, Guralnik JM, Zuliani G, Ferrucci L. Role of muscle mass and muscle quality in the association between diabetes and gait speed. *Diabetes Care.* 2012; 35: 1672-1679

Rodríguez-Mañas L, Fried LP. Frailty in the clinical scenario. *Lancet.* 2015;385: 7-9.

Cadore EL, Casas-Herrero A, Zambom-Ferraresi F, et al. Multicomponent exercises including muscle power training enhance muscle mass, power output, and functional outcomes in institutionalized frail nonagenarians. *Age (Dordr)* 2013;36:773-785.

Cadore EL, Moneo AB, Mensat MM, et al. Positive effects of resistance training in frail elderly patients with dementia after long-term physical restraint. *Age (Dordr)* 2014;36:801-811.

Morley JE, Velas B, van Kan GA, et al. Frailty consensus: A call to action. *J Am Med Dir Assoc.* 2013;14:392-397.

Cadore EL, Rodríguez-Mañas L, Sinclair A, Izquierdo M. Effects of different exercise interventions on risk of falls, gait ability, and balance in physically frail older adults: A systematic review. *Rejuvenation Res.* 2013;16:105-114.

Pahor M, Guralnik JM, Ambrosius WT, Blair S, Bonds DE, Church TS, Espeland MA, Fielding RA, Gill TM, Groessl EJ, King AC, Kritchevsky SB, Manini TM, McDermott MM, Miller ME, Newman AB, Rejeski WJ, Sink KM, Williamson JD; LIFE study investigators. Effect of structured physical activity on prevention of major mobility disability in older adults: the LIFE study randomized clinical trial. *JAMA.* 2014 Jun 18;311(23):2387-96.

Sinclair AJ, Dunning T, Rodriguez-Mañas L. Diabetes mellitus in older people. New insights and residual challenges. *Lancet Diabetes Endocrinol.* 2015;3:275e285.

ADA. Standards of medical care in diabetes. *Diabetes Care.* 2011; 34: 11-61.

Umpierre D, Ribeiro PA, Kramer CK, Leitao CB, Zucatti AT, Azevedo MJ, Gross JL, Ribeiro JP, Schaan BD. Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA.* 2011; 305: 1790-1799.

Hovanec N, Sawant A, Overend TJ, Petrella RJ, Vandervoort AA. Resistance training and older adults with type 2 diabetes mellitus: strength of the evidence. *J Aging Res.* 2012; 284635

Ibañez J, Izquierdo M, Argüelles I, Forga L, Larrión JL, García-Unciti M, Idoate F, Gorostiaga EM. (2005) Twice-weekly progressive resistance training decreases abdominal fat and improves insulin sensitivity in older men with type 2 diabetes. *Diabetes Care.* 2005 Mar;28(3):662-7.

Pitkälä K, Savikko N, Poysti M, Strandberg T, Laakkonen M-L. Efficacy of physical exercise intervention on mobility and physical functioning in older people with dementia: A systematic review. *Experimental Gerontology.* 2013; 48: 85-93.

Izquierdo M, Rodriguez-Mañas L, Sinclair AJ. (2016) What Is New in Exercise Regimes for Frail Older People - How Does the Erasmus Vivifrail Project Take Us Forward? *J Nutr Health Aging.* 2016;20(7):736-7.

Izquierdo M, Rodriguez-Mañas L, Casas-Herrero A, Martínez-Velilla N, Cadore EL, Sinclair AJ. Is It Ethical Not to Prescribe Physical Activity for the Elderly Frail? *J Am Med Dir Assoc.* 2016 Sep 1;17(9):779-81.

Management of Multimorbidities

Fisher K, Griffith L, Gruneir A, Panjwani D, Gandhi S, Sheng LL, et al. Comorbidity and its relationship with health service use and cost in community-living older adults with diabetes: A population-based study in Ontario, Canada. *Diabetes research and clinical practice.* 2016;122:113-23. doi: 10.1016/j.diabres.2016.10.009. PubMed PMID: 27833049.

Kim HM, Seong JM, Kim J. Risk of hospitalization for hypoglycemia among older Korean people with diabetes mellitus: Interactions between treatment modalities and comorbidities. *Medicine (Baltimore).* 2016;95(42):e5016. doi: 10.1097/MD.0000000000005016. PubMed PMID: 27759630; PubMed Central PMCID: PMC5079314.

Pijpers E, Ferreira I, de Jongh RT, Deeg DJ, Lips P, Stehouwer CD, et al. Older individuals with diabetes have an increased risk of recurrent falls: analysis of potential mediating factors: the Longitudinal Aging Study Amsterdam. *Age Aging.* 2012;41(3):358-65. doi: 10.1093/aging/afr145. PubMed PMID: 22156559.

- Bourdel-Marchasson I, Helmer C, Barberger-Gateau P, Peuchant E, Fevrier B, Ritchie K, et al. Characteristics of undiagnosed diabetes in community-dwelling French elderly: the 3C study. *Diabetes research and clinical practice*. 2007;76(2):257-64. PubMed PMID: 17000022.
- Bourdel-Marchasson I, Dubroca B, Manciet G, Decamps A, Emeriau JP, Dartigues JF. Prevalence of diabetes and effect on quality of life in older French living in the community: the PAQUID Epidemiological Survey. *Journal of the American Geriatrics Society*. 1997;45(3):295-301. PubMed PMID: 9063274.
- Pornet C, Bourdel-Marchasson I, Lecomte P, Eschwege E, Romon I, Fosse S, et al. Trends in the quality of care for elderly people with type 2 diabetes: The need for improvements in safety and quality (the 2001 and 2007 Entré Surveys). *Diabetes & metabolism*. 2011;37:152-61. PubMed PMID: 21435929.
- Bongaerts BW, Rathmann W, Heier M, Kowall B, Herder C, Stockl D, et al. Older subjects with diabetes and prediabetes are frequently unaware of having distal sensorimotor polyneuropathy: the KORA F4 study. *Diabetes Care*. 2013;36(5):1141-6. doi: 10.2337/dc12-0744. PubMed PMID: 23275355; PubMed Central PMCID: PMC3631873.
- Sinclair AJ, Bayer AJ, Girling AJ, Woodhouse KW. Older adults, diabetes mellitus and visual acuity: a community-based case-control study. *Age Aging*. 2000;29(4):335-9. PubMed PMID: 10985443.
- Bourdel-Marchasson I, Druet C, Helmer C, Eschwege E, Lecomte P, Le-Goff M, et al. Correlates of health-related quality of life in French people with type 2 diabetes. *Diabetes research and clinical practice*. 2013;101(2):226-35. doi: 10.1016/j.diabres.2013.05.011. PubMed PMID: 23831112.
- Bruderer SG, Bodmer M, Jick SS, Bader G, Schlienger RG, Meier CR. Incidence of and risk factors for severe hypoglycemia in treated type 2 diabetes mellitus patients in the UK--a nested case-control analysis. *Diabetes Obes Metab*. 2014;16(9):801-11. doi: 10.1111/dom.12282. PubMed PMID: 24612200.
- Wang CP, Lorenzo C, Habib SL, Jo B, Espinoza SE. Differential effects of metformin on age related comorbidities in older men with type 2 diabetes. *J Diabetes Complications*. 2017;31(4):679-86. doi: 10.1016/j.jdiacomp.2017.01.013. PubMed PMID: 28190681.
- Chiles NS, Phillips CL, Volpatto S, Bandinelli S, Ferrucci L, Guralnik JM, et al. Diabetes, peripheral neuropathy, and lower-extremity function. *J Diabetes Complications*. 2014;28(1):91-5. doi: 10.1016/j.jdiacomp.2013.08.007. PubMed PMID: 24120281; PubMed Central PMCID: PMC3987662.
- Pearson-Stuttard J, Blundell S, Harris T, Cook DG, Critchley J. Diabetes and infection: assessing the association with glycaemic control in population-based studies. *Lancet Diabetes Endocrinol*. 2016;4(2):148-58. doi: 10.1016/S2213-8587(15)00379-4. PubMed PMID: 26656292.
- Brunström M, Carlberg B. Effect of antihypertensive treatment at different blood pressure levels in patients with diabetes mellitus: systematic review and meta-analyses. *BMJ* 2016; 352:i717.
- Remonti LR, Dias S, Leitão CB, Kramer CK, Klassman LP, Welton NJ, Ades AE, Gross JL. Classes of Antihypertensive Agents and Mortality in Hypertensive Patients with type 2 Diabetes – network meta-analysis of randomized trials. *J Diabetes complications*. 2016; 30: 1192-200.
- Neil HAW, DeMicco DA, Luo D, et al. Analysis of Efficacy and Safety in Patients Aged 65–75 Years at Randomisation. *Diabetes Care*. 2006; 29: 2378-84.
- Ginsberg HN, Elam MB, Lovato LC, et al. Effects of combination lipid therapy in type 2 diabetes mellitus. *N Engl J Med*. 2010; 362: 1563-74.
- Boden WE, Probstfield JL, Anderson T, et al. Niacin in patients with low HDL cholesterol levels receiving intensive statin therapy. *N Engl J Med*. 2011; 365: 2255-67.
- Cheng G, Huang C, Deng H, Wang H. Diabetes as a risk factor for dementia and mild cognitive impairment: a meta-analysis of longitudinal studies. *Intern Med J*. 2012;42(5):484-91. doi: 10.1111/j.1445-5994.2012.02758.x. PubMed PMID: 22372522.
- Sinclair AJ, Girling AJ, Bayer AJ. Cognitive dysfunction in older subjects with diabetes mellitus: impact on diabetes self-management and use of care services. All Wales Research into Elderly (AWARE) Study. *Diabetes research and clinical practice*. 2000;50(3):203-12. PubMed PMID: 11106835.
- Romon I, Rey G, Mandereau-Bruno L, Weill A, Jouglé E, Eschwege E, et al. The excess mortality related to cardiovascular diseases and cancer among adults pharmacologically treated for diabetes--the 2001-2006 ENTRED cohort. *Diabet Med*. 2014;31(8):946-53. doi: 10.1111/dme.12435. PubMed PMID: 24627972.
- Zanders MM, van Steenbergen LN, Haak HR, Rutten HJ, Pruijt JF, Poortmans PM, et al. Diminishing differences in treatment between patients with colorectal cancer with and without diabetes: a population-based study. *Diabet Med*. 2013;30(10):1181-8. doi: 10.1111/dme.12253. PubMed PMID: 23758334.
- Giovannucci E, Harlan DM, Archer MC, Bergenfelz RM, Gapstur SM, Habel LA, et al. Diabetes and cancer: a consensus report. *Diabetes Care*. 2010;33(7):1674-85. doi: 10.2337/dc10-0666. PubMed PMID: 20587728; PubMed Central PMCID: PMC2890380.
- Turnbull PJ, Sinclair AJ. Evaluation of nutritional status and its relationship with functional status in older citizens with diabetes mellitus using the mini nutritional assessment (MNA) tool--a preliminary investigation. *J Nutr Health Aging*. 2002;6(3):185-9. PubMed PMID: 11887244.
- Sanz-Paris A, Gomez-Candela C, Martin-Palmero A, Garcia-Almeida JM, Burgos-Pelaez R, Matia-Martin P, et al. Application of the new ESPEN definition of malnutrition in geriatric diabetic patients during hospitalization: A multicentric study. *Clin Nutr*. 2016;35(6):1564-7. doi: 10.1016/j.clnu.2016.02.018. PubMed PMID: 26997334.
- Ekerstad N, Karlsson BW, Ivanoff SD, et al. Is the acute care of frail elderly patients in a comprehensive geriatric assessment unit superior to conventional acute medical care? *Clin Interv Aging*. 2017; 12: 1-9.
- Meid AD, Lampert A, Burnett A, et al. The impact of pharmaceutical care interventions for medication underuse in older people: a systematic review and meta-analysis. *Br J Clin Pharmacol*. 2015; 80: 768-776.

Hypoglycemia Management

- ADVANCE Collaborative Group, Patel A, MacMahon S, et al. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2008;358:2560-72.
- Action to Control Cardiovascular Risk in Diabetes Study Group, Gerstein HC, Miller ME, et al. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545-59.
- Duckworth W, Abraira C, Moritz T, et al. Glucose control and vascular complications in veterans with type 2 diabetes. VDAT Study. *N Engl J Med* 2009;360:129-39.
- Bremer JP, Jauch-Chara K, Hallschmid M, Schmid S, Schultes B. Hypoglycemia unawareness in older compared with middle-aged patients with type 2 diabetes. *Diabetes Care*. 2009;32:1513-7.

Shorr RI, Ray WA, Daugherty JR, et al. Incidence and risk factors for serious hypoglycemia in older persons using insulin or sulphonylurea. *Arch Int Med* 1997; 157: 1681-6.

Yaffe K, Falvey CM, Hamilton N, et al. Association between hypoglycaemia and dementia in a biracial cohort of older adults with diabetes mellitus. *JAMA Intern Med* 2013;173:1300-6.

Feinkohl I, Aung PP, Keller M, et al.; Edinburgh Type 2 Diabetes Study (ET2DS) Investigators. Severe hypoglycaemia and cognitive decline in older people with type 2 diabetes: the Edinburgh type 2 diabetes study. *Diabetes Care* 2014;37:507-15.

Malanda UL, Welschen LMC, Riphagen II, et al. Self-monitoring of blood glucose in patients with type 2 diabetes mellitus who are not using insulin. *Cochrane Database Syst Rev* 2012;1:CD005060.

Abdelhafiz AH, Sinclair AJ. Hypoglycemia in residential care homes. *Br J Gen Pract*. 2009 Jan 1; 59: 49-50.

Munshi, MD, Slyne C, Segal AR, Saul N, Lyons C, Weinger K. Liberating A1C goals in older adults may not protect against the risk of hypoglycemia. *J Diabetes Complications*. DOI:<http://dx.doi.org/10.1016/j.jdiacomp.2017.02.014>.

Thorpe CT, Gellad WF, Good CB, Zhang S, Zhao X, Mor M, Fine1 MJ. Tight Glycemic Control and Use of Hypoglycemic Medications in Older Veterans With Type 2 Diabetes and Comorbid Dementia. *Diabetes Care* 2015; 38: 588-95.

Greco D, Pisciotta M, Gambina F, et al. Severe hypoglycemia leading to hospital admission in type 2 diabetic patients aged 80 years or older. *Exp Clin Endocrinol Diabetes* 2010;118:215-9.

McCoy RG, Lipska KJ, Yao X, et al. Intensive Treatment and Severe Hypoglycemia Among Adults With Type 2 Diabetes. *JAMA Intern Med*. 2016; 176: 969-78.

Sussman JB, Kerr EA, Saini SD, et al. Rates of Deintensification of Blood Pressure and Glycemic Medication Treatment Based on Levels of Control and Life Expectancy in Older Patients With Diabetes Mellitus. *JAMA Intern Med* 2015;175:1942-9.

Munshi MN, Slyne C, Segal AR, et al. Simplification of Insulin Regimen in Older Adults and Risk of Hypoglycemia. *JAMA Intern Med* 2016; 176: 1023-5.

Sjöblom P, Tengblad A, Lofgren UB et al. Can diabetes medication be reduced in elderly patients? An observational study of diabetes drug withdrawal in nursing home patients with tight glycaemic control. *Dib Res Clin Pract* 2008; 82: 197-202.

Abdelhafiz AH, Chakravorty P, Gupta S et al. [Can hypoglycaemic medications be withdrawn in older people with type 2 diabetes?](#) *Int J Clin Pract* 2014; 68: 790-2.

Caverly TJ, Fagerlin A, Zikmund-Fisher BJ, et al. Appropriate Prescribing for Patients With Diabetes at High Risk for Hypoglycemia: National Survey of Veterans Affairs Health Care Professionals. *JAMA Intern Med* 2015; 175: 1994-6.

Sinclair A, Armes DG, Randhawa G, et al. Caring for older adults with diabetes mellitus: characteristics of carers and their prime roles and responsibilities. *Diabet Med*. 2010;27:1055-9.

Sinclair AJ, Gadsby R, Hillson R, et al. Brief report: Use of the Mini-Cog as a screening tool for cognitive impairment in diabetes in primary care. *Diabetes*

Primary and Community Care

Bethel MA, Sloan FA, Belsky D, Feinglos MN. Longitudinal incidence and prevalence of adverse outcomes of diabetes mellitus in elderly patients. *Arch Intern Med* 2007; 167 (9): 921-7.

Kirkman MS, Briscoe VJ, Clark N, Florez H, Haas LB, Halter JB, et al. Diabetes in older adults: a consensus report. *J Am Geriatr Soc* 2012; 60 (12): 2342-56.

Lerberghe WV. The world health report 2008 [electronic resource]. Primary health care : now more than ever. Geneva: WHO, 2008.

Groenewegen PP, Dourgnon P, Gress S, Jurgutis A, Willems S. Strengthening weak primary care systems: steps towards stronger primary care in selected Western and Eastern European countries. *Health Policy* 2013; 113 (1-2): 170-9.

Boeckxstaens P, De Graaf P. Primary care and care for older persons: position paper of the European Forum for Primary Care. *Qual Primary Care* 2011; 19 (6): 369-89.

Hoskins PL. Sharing the care of diabetic patients between hospital and general practitioners: does it work? *Diabetic Med* 1993; 10 (1): 81.

Griffin S. Diabetes care in general practice: meta-analysis of randomised control trials. *Br Med J* 1998; 317 (7155): 390-6.

Diabetes Integrated Care Evaluation T. Integrated Care For Diabetes: Clinical, Psychosocial, and Economic Evaluation. *Br Med J* 1994; 308 (6938): 1208-12.

Hayes TM, Harries J. Randomised controlled trial of routine hospital clinic care versus routine general practice care for type II diabetics. *Br Med J (Clin Res Edn)* 1984; 289 (6447): 728-30.

Rodríguez-Mañas L, Bayer AJ, Kelly M, Zeyfang A, Izquierdo M, Laosa O, Hardman TC, Sinclair AJ, Moreira S, Cook J . An evaluation of the effectiveness of a multi-modal intervention in frail and pre-frail older people with type 2 diabetes--the MID-Frail study: study protocol for a randomised controlled trial.; MID-Frail Consortium. *Trials*. 2014 Jan 24;15:34.

Kennedy M. Primary and Community Care of Diabetes in Older People. In: *Diabetes in Old Age*; Eds: Sinclair AJ, Dunning T, Rodriguez-Manas L, Munshi M; 4th edition, John Wiley & Sons, Chichester, UK 2017; pp 376-394

Sinclair AJ, Gadsby R, Penfold S, Croxson SC, Bayer AJ. Prevalence of diabetes in care home residents. *Diabetes Care* 2001; 24 (6): 1066-8.

Sinclair AJ, Task & Finish Group of Diabetes UK. Good clinical practice guidelines for care home residents with diabetes: an executive summary. *Diabetic Med* 2011; 28 (7): 772-7.

Benetos A, Novella J-L, Guerci B, Bickle J-F, Boivin J-M, Cuny P, et al. Pragmatic diabetes management in nursing homes: individual care plan. *J Am Medical Directors Assoc* 2013; 14 (11): 791-800.

Sinclair AJ[javascript:void\(0\);](#), Abdelhafiz A, Rodríguez-Mañas L. Frailty and Sarcopenia - newly emerging and high impact complications of diabetes. *J Diab Complications* 2017; *In press*

Sinclair AJ, Paolisso G, Castro M, Bourdel-Marchasson I, Gadsby R, Rodriguez Manas L. European Diabetes Working Party for Older People 2011 Clinical Guidelines for Type 2 Diabetes Mellitus. Executive Summary – A Report of the European Diabetes Working Party for Older People (EDWPOP) Revision Group on Clinical Practice Guidelines for Type 2 Diabetes Mellitus. *Diabetes Metab* 2011; 37: S27-38.

Inpatient Care

Schneider ALC, Kalyani RR, Golden S, et al. Diabetes and prediabetes and risk of hospitalization: the Atherosclerosis Risk in Communities (ARIC) Study. *Diabetes Care* 2016; 39: 772-9.

van den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med* 2001; 345: 1359-67.

Cook CB, Kongable GL, Potter DJ, et al. Inpatient glucose control: a glycemic survey of 126 U.S. hospitals. *J Hosp Med* 2009; 4: E7-E14.

Falciglia M, Freyberg RW, Almenoff PL, et al. Hyperglycemia related mortality in critically ill patients varies with admission diagnosis. *Crit Care Med* 2009; 37: 3001-9.

Kagansky N, Levy S, Rimon E, et al. Hypoglycemia as a predictor of mortality in hospitalized elderly patients. *Arch Intern Med* 2003; 163: 1825-9.

Cox DJ, Kovatchev BP, Gonder-Frederick LA, et al. Relationships between hyperglycemia and cognitive performance among adults with type 1 and type 2 diabetes. *Diabetes Care* 2005; 28: 71-7.

Umpierrez GE, Smiley D, Jacobs S, et al. Randomized study of basal-bolus insulin therapy in the inpatient management of patients with type 2 diabetes undergoing general surgery (RABBIT 2 surgery). *Diabetes Care* 2011; 34: 256-61.

Pasquel FJ, Gianchandani R, Rubin DJ, et al. Efficacy of sitagliptin for the hospital management of general medicine and surgery patients with type 2 diabetes (Sita-Hospital): a multicentre, prospective, open-label, non-inferiority randomised trial. *Lancet Diabetes Endocrinol* 2017; 5: 125-33.

Chiang R, Fairclough E, Tan E, Abdelhafiz A H. Effect of Diabetes Service for Older People on Length of Hospital Stay. *J Am Geriatr Soc* 2008; 56: 2143-5.

Wilson C, Curtis J, Lipke S, et al. Nurse case manager effectiveness and case load in a large clinical practice: implications for workforce development. *Diabet Med* 2005; 22: 1116-20.

Worrall G, Knight J. Continuity of care is good for elderly people with diabetes: Retrospective cohort study of mortality and hospitalization. *Can Fam Physician*

<http://www.ncbi.nlm.nih.gov/pubmed/21252120> - #2011; 57: e16-e20.

Educational Aspects for Health Professionals in Clinical Settings

Fisher J. Frail Fail: Four thoughts on framing frailty teaching. Available from:
<https://britishgeriatricsociety.wordpress.com/2014/06/24/frail-fail-thouhts-on-frailty-teaching> (Accessed 1st March 2017).

Module-Frailty. Camden Clinical commissioning Group. Available from: <https://gps.camdnccg.nhs.uk/education-module-frailty> (Accessed 1st March 2017).

de Villiers Frailty Continuing Medical education 2013 available at:
<http://www.cmej.org.za/index.php/cmej/article/view/2868/3235> (last accessed January 31st 2017).

Diener E, Biswas-Diener R. Flourishing Scale 2009. Available from
<https://internal.psychology.illinois.edu/~ediener/Documents/FS.pdf> (Accessed 23rd February 2017).

Currie K, Spink J and Rajendran M. Well-written health information: A guide. Communicating with Consumers series volume 1. 2000, Victorian Government Department of Human Services, Melbourne. Available from:
<http://www.health.vic.gov.au/consumer/pubs/written.htm> (Accessed 30th December 2016).

Renteln-Kruse W, Minder C. It is time to detect preclinical signs of incipient frailty. *Journal American Geriatrics Society*. 2016; 64(10):2166-67.

Kirk E, Paddon-Jones D. Protecting muscle mass and function in older adults during bed rest. *Current Opinion Clinical Nutrition Metabolic Care*. 2010; 13(1):34-39.

Persz-Zepeda M, Cesari M, Grrcia-Pena C. Predictive value of frailty indices for adverse outcome in older adults. *Rev Inves clin*. 2016;68:92-98.

Chan D-C, Tsou H-H, Chang C-B, Yang R-S, Tsauo J-Y, Chen C-Y, Hsiao C-F, Hsu Y-T, Chen C-H, Chang S-F, Hsiung C-A,kou K. Integrated care for geriatric frailty and sarcopenia: a randomized control trial. *Journal Cachexia, Sarcopenia and Muscle*. 2017; 8:78-88.

de Labra C, Guimaraes-Pinheiro C, Maseda A, Lorenzo T, Millan-Calenti J. Effects of physical exercise interventions in frail older adults: a systematic review of randomized control trials. *BMC Geriatrician*. 2005;15: 154 doi: 10.1186/s12877-015-0144-4.

Sinclair A, Conroy S, Bayer A.: Impact of diabetes on physical function in older people. *Diabetes Care*. 2008, 31: 233-235

Keough M, Field T, Gurwitz J. A model of community-based interdisciplinary team training in the care of the frail elderly. *Academic Medicine*. 2002; 7(9):936

Dyer C, Feldt K, Lindemann D, Busby-Whitehead J, Greenberg S, Kennedy R, Flaherty E. Frail older patient care by interdisciplinary teams: a primer for generalists. *Gerontology Geriatric Education*. 2003;24(2): 51-62.

Hoogendoijk E, van Hout H, Heymans M, van der Horst H, Frijters D, Broese van Groenou M, Deeg D, Huisman M. Explaining the association between educational level and frailty in older adults: results of a 13-year longitudinal study in the Netherlands. *Annals of Epidemiology*. 2014;4(7):538-544.

Glisky E, Rubin S, Davidson S. Source of memory in older adults: an encoding and retrieving problem. *Journal Experimental Psychology*. 2001;27(5):1131-1146.

Brand G, Carr S, Etherton-Beer C. Depth of field: teaching the art of reflection to medical students. *Journal Gerontology and Geriatric Research*. 2016;5(6):7182.

Brand G, Osborne A, Carroll M, Carr S, Etherton-Beer C. Do photographs, older adults' narratives and collaborative dialogue foster anticipatory reflection (preflection) in medical students? *BMC Medical Education*. 2016;16: 289 doi: [10.1186/s12909-016-0802-2](https://doi.org/10.1186/s12909-016-0802-2)

Graham I, Logan J, Harrison M, Straus S, Tetroe J, Caswell W, Robinson N: Lost in knowledge translation: time for a map? *Journal Continuing Education Health Professionals*. 2006;26: 13-24. 10.1002/chp.47.

Research Directions

Morley JE, Malmstrom TK, Rodriguez-Mañas L, Sinclair AJ. Frailty, sarcopenia and diabetes. *J Am Med Dir Assoc*. 2014; 15: 853-9.

Kalyani RR, Corriere M, Ferrucci L. Age-related and disease-related muscle loss: the effect of diabetes, obesity, and other diseases. *Lancet Diabetes Endocrinol*. 2014; 2: 819-29.

Sinclair A, Dunning T, Rodriguez-Mañas L. Diabetes in older people: new insights and remaining challenges. *Lancet Diabetes Endocrinol*. 2015; 3: 275-85.

- Sinclair AJ, Paolisso G, Castro M, Bourdel-Marchasson I, Gadsby R, Rodriguez Mañas L; European Diabetes Working Party for Older People. European Diabetes Working Party for Older People 2011 clinical guidelines for type 2 diabetes mellitus. Executive summary. *Diabetes Metab.* 2011; 37 Suppl 3: S27-38.
- Kirkman SM, Briscoe VJ, Clark N, Florez H, Haas LB, Harker JB et al. Diabetes in older adults: a consensus report. *J Am Geriatr Soc* 2012; 60: 2342-56
- Munshi MN, Florez H, Huang ES, Kalyani RR, Mupanomunda M, Pandya N, et al. Management of Diabetes in Long-term Care and Skilled Nursing Facilities: A Position Statement of the American Diabetes Association. *Diabetes Care.* 2016; 39: 308-18.
- Soriguer F, Goday A, Bosch-Comas A, Bordiú E, Calle-Pascual A, Carmena R, et al. Prevalence of diabetes mellitus and impaired glucose regulation in Spain: the Diabetologia 2012; 55: 88-93.
- Castro-Rodriguez M, Carnicero JA, Garcia-Garcia FJ, Walter S, Morley JE, Rodriguez-Artalejo F, et al. Frailty as a Major Factor in the Increased Risk of Death and Disability in Older People With Diabetes. *J Am Med Dir Assoc.* 2016; 17: 949-55.
- García-Esquinas E, Graciani A, Guallar-Castillón P, López-García E, Rodríguez-Mañas L, Rodríguez-Artalejo F. Diabetes and risk of frailty and its potential mechanisms: a prospective cohort study of older adults. *J Am Med Dir Assoc.* 2015; 16: 748-54.
- Monette MC, Baird A, Jackson DL. A meta-analysis of cognitive functioning in nondemented adults with type 2 diabetes mellitus. *Can J Diabetes.* 2014; 38: 401-8.
- Speight J, Sinclair AJ, Browne JL, Woodcock A, Bradley C. Assessing the impact of diabetes on the quality of life of older adults living in a care home: validation of the ADDQoL Senior. *Diabet Med.* 2013; 30: 74-80.
- Rodríguez-Sánchez B, Angelini V, Feenstra T, Alessie RJ. Diabetes-Associated Factors as Predictors of Nursing Home Admission and Costs in the Elderly Across Europe. *J Am Med Dir Assoc.* 2017; 18: 74-82.
- Rodríguez-Mañas L, Bayer AJ, Kelly M, Zeyfang A, Izquierdo M, Laosa O, et al.; MID-FRAIL Consortium. An evaluation of the effectiveness of a multi-modal intervention in frail and pre-frail older people with type 2 diabetes--the MID-Frail study: study protocol for a randomised controlled trial. *Trials.* 2014; 15: 34.
- Landi F, Cesari M, Calvani R, Cherubini A, Di Bari M, Bejuit R, et al. The "Sarcopenia and Physical fRAILty IN older people: multi-component Treatment strategies" (SPRINTT) randomized controlled trial: design and methods. *Aging Clin Exp Res.* 2017; 29: 89-100.
- Rodríguez-Mañas L, Rodríguez-Artalejo F, Sinclair AJ. The Third Transition: The Clinical Evolution Oriented to the Contemporary Older Patient. *J Am Med Dir Assoc.* 2017; 18: 8-9.
- Sinclair A, Morley J. Frailty and diabetes. *Lancet* 2013; 382: 1386-7.
- Sinclair AJ. Towards a minimum data set for intervention studies in type 2 diabetes in older people. *J Nutr Health Aging.* 2007; 11: 289-93.
- Sinclair A, Morley JE, Rodriguez-Mañas L, Paolisso G, Bayer T, Zeyfang A. Diabetes mellitus in older people: position statement on behalf of the International Association of Gerontology and Geriatrics (IAGG), the European Diabetes Working Party for Older People (EDWPOP), and the International Task Force of Experts in Diabetes. *J Am Med Dir Assoc.* 2012; 13: 497-502.
- WHO. World report on aging and health. 2015. Available at: <http://www.who.int/ageing/events/world-report-2015-launch/en/>
- Beard JR, Officer A, de Carvalho IA, Sadana R, Pot AM, Michel JP, et al. The World report on aging and health: a policy framework for healthy aging. *Lancet* 2016; 387: 2145-54.